
Your Treatment Guide





What is ZYNLONTA®?

ZYNLONTA® (loncastuximab tesirine-lypl) is a prescription medicine used to treat adults with certain types of large B-cell lymphoma that has come back (relapsed) or that did not respond to previous treatment (refractory), who have already received two or more treatments for their cancer.

It is not known if ZYNLONTA® is safe and effective in children.

The approval of ZYNLONTA® is based on a type of response rate. There is an ongoing study to confirm the clinical benefit of ZYNLONTA®.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of ZYNLONTA®?

ZYNLONTA® may cause serious side effects, including:

 Fluid retention. Your body may hold too much fluid during treatment with ZYNLONTA[®]. This can be serious. Tell your healthcare provider if you develop new or worsening swelling or puffiness, weight gain, chest pain, shortness of breath, or trouble breathing

Please see additional Important Safety Information throughout this brochure, and full **Prescribing Information** for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.







2

Topics

Ask Your Doctor If 3 Staying Up To Date With 15 **ZYNLONTA®** Is Right For You **ZYNLONTA® Questions To Ask Your Doctor** 16 **ZYNLONTA®** Specifically **About ZYNLONTA® Targets Cancer Caregiver Support** What ZYNLONTA® Infusions 5 20 & Tips **Are Like Track Your Results To Expect With** 7 22 **Treatment Progress ZYNLONTA® Important Safety** 23 9 **Managing Possible Side** Information **Effects** Resources **ADVANCING Patient Support** 13 25

Click topics below to visit the page >

Please see additional Important Safety Information throughout this brochure, and full **Prescribing Information** for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.





ZYNLONTA® is a targeted therapy for adults who have tried 2 or more therapies before, but the DLBCL came back (relapsed), or those therapies did not work (refractory).

ZYNLONTA®:



Is not a traditional chemotherapy _____ Go to page 4 > Specifically targets DLBCL cancer cells Is given every 3 weeks _____ Go to page 5 > Was studied in adults who had received a variety of DLBCL therapies _____ Go to page 7 >

IMPORTANT SAFETY INFORMATION (continued)

- Low blood cell counts (platelets, red blood cells, and white blood cells). Low blood cell counts are common with ZYNLONTA® but can also be serious or severe. Your healthcare provider will monitor your blood counts during treatment with ZYNLONTA®. Tell your healthcare provider right away if you get a fever of 100.4°F (38°C) or above, or any bruising or bleeding
- Infections. Serious infections, including infections that can cause death, have happened in people treated with ZYNLONTA®. Tell your healthcare provider right away if you have new or worsening signs or symptoms of infection

Please see additional Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on ZYNLONTA.com.













Not an actual patient.

ZYNLONTA® uses a find, bind, and release strategy to target cancer cells

ZYNLONTA® is what's known as an antibody-drug conjugate (or ADC). It is different than traditional **chemotherapy**.

ZYNLONTA® contains an **antibody**, (a protein that works with your immune system), that is attached to an anti-cancer drug.





The antibody finds and binds to a specific area of the cancer cell called CD19

The cancer cell absorbs **ZYNLONTA®**

Once inside, ZYNLONTA® breaks apart and the anti-cancer drug is released

This causes the cancer cell to die

IMPORTANT SAFETY INFORMATION (continued)

• Skin reactions. Serious skin reactions have happened in people treated with ZYNLONTA®. Tell your healthcare provider if you get new or worsening skin reactions, including sensitivity to sunlight, skin rash, peeling, redness or irritation. You may burn more easily or get severe sunburns

Please see additional Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.





ZYNLONTA® lets you focus on the time between treatments





Not an actual patient.


ZYNLONTA® is given at a clinic or infusion center

ZYNLONTA® is given:

Through an **intravenous infusion** (given via a needle placed in the vein)

In about 30 minutes

Once every 3 weeks



Your doctor will decide how many treatments you need.

Before each treatment, you may need a prescription steroid (dexamethasone) to lower your chance of side effects. Go to next page for more information

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of ZYNLONTA® include:

- · feeling tired or weak
- skin rash
- swelling
- nausea
- muscle or joint pain
- increase in blood sugar (hyperglycemia)
- changes in certain blood or laboratory tests

ZYNLONTA® may cause fertility problems in males which may affect your ability to father children. Talk to your healthcare provider if this is a concern for you. These are not all of the possible side effects of ZYNLONTA®.



Please see additional Important Safety Information throughout this brochure, and full **Prescribing Information** for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.





Learn the facts about ZYNLONTA® treatment



Not an actual patient.

Your doctor may give you a steroid (dexamethasone) before ZYNLONTA®

2 times a day



for 3 days



starting the day before infusion(s)

If you have severe side effects, your doctor may:







Stop your treatment

Delay your treatment

Change your dose of ZYNLONTA®

What to do if you miss a dose



Alert your doctor's office right away to talk about your options

- Your doctor should do blood tests regularly to check for side effects of ZYNLONTA®
- You will be monitored for infusion site reactions, infections, and skin reactions. You may also need CT scans to monitor the cancer

IMPORTANT SAFETY INFORMATION (continued)

What should I avoid while receiving ZYNLONTA®?

Avoid or limit your exposure to sunlight, including sunlight through glass, such as buildings or vehicle windows and artificial sunlight such as sunlamps or tanning beds. Exposure to sunlight during treatment with ZYNLONTA® can cause skin reaction or rash. Use sun protection measures such as sunscreen and wear loose-fitting clothes that cover your skin while out in sunlight.

Please see additional Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.









The participants had past relapsed/refractory (R/R) DLBCL therapies



Chemotherapy



RITUXAN® (rituximab)



CAR-T therapy



The trial included 145 people with R/R DLBCL

- 88% had a type of disease called DLBCL not otherwise specified (NOS)
- 20% had DLBCL arising from slow-growing lymphoma
- 7% had fast-growing B-cell lymphoma

Age range

- 23 to 94
- 66 was the median age

Sex

- Male: 59%
- Female: 41%

Race

- Caucasian: 90%
- Other: 10%

Past therapies

- At least 2, and as many as 7 therapies
- 17% had previous stem cell transplant
- 9% had previous CAR-T-cell therapy
- More than half of patients (63%) did not respond to their previous treatment

After the trial

 68 people who received ZYNLONTA® went on to other therapies

IMPORTANT SAFETY INFORMATION (continued)

Before you receive ZYNLONTA®, tell your healthcare provider about all of your medical conditions, including if you:

- · have an active infection or have had one recently
- have liver problems
- are pregnant or plan to become pregnant. ZYNLONTA® can harm your unborn baby
- are breastfeeding or plan to breastfeed. It is not known if ZYNLONTA® passes into breast milk. Do not breastfeed during treatment with ZYNLONTA® and for 3 months after the last dose of ZYNLONTA®



Please see additional Important Safety Information throughout this brochure, and full **Prescribing Information** for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.





In the clinical trial that led to ZYNLONTA® being approved for use,

Quick and effective results were seen in some people with ZYNLONTA®



Not an actual patient.



24%

had all signs of the cancer go away (a "complete response") 35 out of 145

24% had some of the cancer lessen (a "partial response") 35 out of 145

Of those who responded:



Half of people saw results after about **1 month** or 2 treatments*



Half of people saw results last for about 10 months[†]

The approval of ZYNLONTA® is based on a type of response rate. There is an ongoing study to see how well the drug works

Partial response: Signs of cancer have been reduced.

Complete response: Signs of cancer have disappeared. This does not always mean the cancer has been cured.


Overall response: Includes both complete responses and partial responses.

IMPORTANT SAFETY INFORMATION (continued)

Females who can become pregnant:

- your healthcare provider may do a pregnancy test before starting treatment with ZYNLONTA®
- you should use effective birth control (contraception) during treatment with ZYNLONTA® and for 10 months after the last dose
 of ZYNLONTA®. Talk to your healthcare provider about effective birth control. Tell your healthcare provider right away if you
 become pregnant or think that you are pregnant during treatment with ZYNLONTA®

Please see additional Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.



[†]This is called the "median duration of response." That means half of people's cancer kept on responding to treatment without it growing or spreading for **10.3 months** and the other half of people's cancer responded for a shorter amount of time.











Talk to your doctor's office if you experience serious side effects



Swelling from fluid buildup (fluid retention):

This can be a serious side effect.

Tell your doctor's office if you have new or worsening:

- Swelling or puffiness
- · Weight gain
- Chest pain
- Shortness of breath
- Trouble breathing



Low blood cell counts (platelets, red blood cells, and white blood cells):

Low blood cell counts are common with ZYNLONTA®. But that can also be serious or severe. Your doctor will monitor your blood counts during treatment with ZYNLONTA®.

Tell your doctor's office right away if you get a fever of 100.4°F (38°C) or above, or if you have any bruising or bleeding.



Infections:

Serious infections, including infections that can cause death, have happened in individuals treated with ZYNLONTA®.

Tell your doctor's office right away if you have new or worsening signs or symptoms of infection, including:

- Fever
- Chills
- Flu-like symptoms, like cough, tiredness or weakness, and body aches
- Headache
- Breathing problems
- Cuts or scrapes that are red, warm, swollen or painful



Skin reactions:

Serious skin reactions have been reported in individuals treated with ZYNLONTA®.

Tell your doctor's office if you get new or worsening skin reactions, including:

- Sensitivity to sunlight
- A skin rash, peeling, redness, or irritation

Your skin may burn more easily or get severe sunburns.

If you have serious side effects, your doctor may stop or delay your treatment or change the dosage



Please see additional Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.





What to avoid during ZYNLONTA® treatment



Not an actual patient.

The most common side effects of ZYNLONTA® include:

- feeling tired or weak
- skin rash
- swelling
- nausea

- · muscle or joint pain
- increase in blood sugar (hyperglycemia)
- changes in certain blood or laboratory tests

ZYNLONTA® may cause fertility problems in males which may affect your ability to father children. Talk to your healthcare provider if this is a concern for you. These are not all of the possible side effects of ZYNLONTA®. Call your healthcare provider for medical advice about side effects.

If you have certain side effects, your doctor may have you:



Avoid or limit your exposure to sunlight:

This includes sunlight through glass, such as buildings or vehicle windows and artificial sunlight, such as sunlamps or tanning beds. If you must go out in the sun, be sure to wear wide brimmed hats and use high SPF sunscreen.

Exposure to sunlight during treatment with ZYNLONTA® can cause skin reaction or rash.

Use sun protection measures such as sunscreen and wear loose-fitting clothes that cover your skin while out in sunlight.



Before you receive ZYNLONTA®,

Tell your doctor's office about all of your medical conditions, including if you:

- · Have an active infection or have had one recently
- Have liver problems
- Are pregnant or plan to become pregnant because ZYNLONTA® can harm an unborn baby
- Are breastfeeding or plan to breastfeed. It is not known if ZYNLONTA® passes into breast milk. Do not breastfeed during treatment with ZYNLONTA® and for 3 months after the last dose of ZYNLONTA®



Please see additional Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.





Reach out to your doctor about side effects



Not an actual patient.

For females who can become pregnant

If you have certain side effects:

- You should use effective birth control (contraception) during treatment with ZYNLONTA® and for 10 months after the last dose of ZYNLONTA®
- Talk to your doctor's office about effective birth control. Tell your doctor right away if you become pregnant or think that you are
 pregnant during treatment with ZYNLONTA®
- Your doctor's office may do a pregnancy test before starting treatment with ZYNLONTA®

For males with female partners who can become pregnant,

- You should use effective birth control (contraception) during treatment with ZYNLONTA® and for 7 months after the last dose of ZYNLONTA®
- Talk to your doctor's office about effective birth control. Tell your doctor right away if your partner becomes pregnant during your treatment with ZYNLONTA®

Share a list of medicines you take with your doctor's office and pharmacists.

This includes prescription and over-the-counter medicines, vitamins, and herbal supplements







Tips for common side effects



Not an actual patient.

Feeling tired or weak

- Let others help you do things
- · Balance brief, light exercise with short naps, if needed
- · Follow good eating and drinking habits. Drink water, and limit alcohol and caffeine

Skin rash

- Use only mild soaps
- Use a mild nonperfumed lotion to moisturize
- · Use sunscreen and skin-covering clothes when going out
- Shower in lukewarm (not hot) water

Swelling

- Wear loose clothes
- · Prop up your feet when lying or sitting
- Wear compression clothes
- Eat a low-sodium diet. Salt can make swelling worse
- · Your doctor may prescribe a diuretic, which helps your body rid itself of fluid and salt faster

Muscle or joint pain

- Keep track of pain levels and what helps you reduce pain
- Take pain medicine as prescribed. Don't wait until the pain gets to be too much
- · Consider reaching out to a pain specialist
- Ask about alternative therapies, like massage, acupuncture, etc

Nausea

- Take an anti-nausea medicine
- Drink plenty of water and fluids
- Avoid foods that make you feel nauseous
- · Try deep breathing, hypnosis, and other stress-relief routines that help some people with nausea

Increase in blood sugar (hyperglycemia)

You may need to change a few things. Consult your doctor for more info


Changes in certain blood or lab tests


You may need to make a few changes. Your doctor's office will share ways to help

Please see additional Important Safety Information throughout this brochure, and full Prescribing Information for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.











Get personalized help with savings, health insurance support, and treatment details

Help reach your goals with our comprehensive support program, ADVANCING Patient Support. Once enrolled, you can:





Advance to your treatment goals



Overcome common barriers to treatment



Get clear answers to questions

Once enrolled, your dedicated case manager will get to know you and your specific case. You can call them when you need help clarifying complicated information. You can also get answers to questions you might have, like:

- Will insurance cover treatment?
- How much will this cost me?
- How many treatments will be covered?

- Do I qualify for financial assistance?
- Can I talk to someone who knows how ZYNLONTA® works?

Ask your doctor's office about enrolling in ADVANCING Patient Support



Please see Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.





Get clear answers and personalized help



Not an actual patient.

Ask your doctor about how ADVANCING Patient Support could help



Find out if you qualify for:

- **\$0 Copay:** If you're commercially insured and eligible, ADVANCING Patient Support can help your doctor's office enroll you in the copay program. Once you're signed up, you may pay as little as \$0 per treatment*
- Coverage of therapy costs: If you are uninsured or underinsured. ADVANCING Patient Support can help you find out if you qualify for the Patient Assistance Program




Your case manager can help you understand:


- Coverage: Whether or not ZYNLONTA® is covered under your health plan
- Cost: What the cost may be for your therapy
- Necessaries: What, if any, approvals or additional information you need before receiving therapy



Get in touch with a nurse who can:


- Answer questions about ZYNLONTA® and your health
- Prepare you for conversations with your doctor
- Connect you to additional support











Get updates right to your inbox



ADC Therapeutics is committed to providing you with all-important education and treatment information. Our hope is that key information helps drive more productive conversations with your care team.

You'll receive:



Easy-to-understand information about ZYNLONTA®



Questions to drive conversations with your doctors and their staff



More information about ADVANCING Patient Support



Caregiving, from logistics to heart-to-hearts

Go to the ZYNLONTA® Updates Program page on ZYNLONTA.com to sign up >





Have more productive conversations with your doctor







Not an actual patient.



Checklist of Q & A

At this time, discussions with your doctor about your options are crucial. The questions below can help guide the conversation at your next appointment.

Is ZYNLONTA® right for my specific DLBCL?	How well does ZYNLONTA® work?	How is ZYNLONTA® given?
How long does each treatment take? How often is ZYNLONTA® given?	Does it matter what therapy I was on before?	What are possible side effects I should look out for?

Go to Table of Contents > 📙





Have more productive conversations with your doctor (cont'd)



Not an actual patient.

٢	<u>.</u>
	žΞ
ι	

Checklist of Q & A

At this time, discussions with your doctor about your options are crucial. The questions below can help guide the conversation at your next appointment.

When should I contact my office about side effects?	Is there anything I should avoid taking or doing while on ZYNLONTA®?	What are some trusted sources that can help me understand DLBCL?			
Notes					

Go to Table of Contents >





Tracking contact info

Healthcare provide	ers' contact information		
<u>Ω</u> * (P) Oncologist	(Oncologist Office	[Ωˇ ♠] Primary Care
·	Practitioner	Administrator	Physician
Notes			

Please see Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on <u>ZYNLONTA.com</u>.





Additional Notes

Notes	
	-
	-
	G0 10
	lable
	or Cont
	Contents >

Please see Important Safety Information throughout this brochure, and full <u>Prescribing Information</u> for ZYNLONTA®, including Patient Information, on <u>ZYNLONTA.com</u>.



Being prepared can change how appointments go





Not an actual patient.

Caregiver resources

Don't be afraid to ask questions if you don't understand something. Part of the care team's job is to make sure everything is clear to the person you care for and you.



Having open conversations with their doctor can make a big difference. Consider making a list of questions to ask. Some topics you may wish to ask about are:



About their treatment

What should the person you care for be encouraged to do while undergoing treatment?



Planning for the infusion

What is the process like before the infusion?



About your role

How might being a caregiver affect the day-to-day?



Important things to watch out for

What side effects should we contact the office about?

Go to the caregiver section of ZYNLONTA.com for more information >









Helpful caregiver tips



Not an actual patient.

Caregiver resources continued

Providing care is never an easy job. And it's just as important to take care of yourself. Here are some helpful tools and tips to help you be the best caregiver you can be.



Don't be afraid to speak up.

Ask questions and take notes during doctor visits. Your input is important.

Keep them involved in their care.

Let them make their own decisions. You and their care team can help keep things on track.

Also take care of yourself.

Meditate, exercise, get good rest, see your friends, take some time for yourself... do whatever you need to be at your best.

Education helps.

Learning more can help you assist your loved one during doctor visits. It can also help you relate to your loved one's experience.

You're not alone.

Taking advantage of caregiver resources—including online support groups—can give you an outlet.

1-202-454-3970

1-800-445-8106

1-800-500-9976

Other resources

Caregiver Action Network
Family Caregiver Alliance
Friends' Health Connection
Lymphoma Research Foundation Helpline
Lymphoma Peer Support Network
Education Programs

www.caregiveraction.org
www.caregiver.org
www.friendshealthconnection.org
helpline@lymphoma.org

https://lymphoma.org/resources/supportservices/lsn/

https://lymphoma.org/resources/educationresources/programs



Please see Important Safety Information throughout this brochure, and full **Prescribing Information** for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.



ZYNLONTA® may be able to

Defining your own "here" and getting there

The state of the s	A
	A

	Where do you want to be? Close your eyes for a moment and think about how you'd fill this in:				ent	help you get you to your "here," wherever			
Treatment goal setting	" most like myself."		is my 'here,' a place I feel				that may be.		
Notes or questions									
to ask your care team		:	/ /	:		:		:	
Infusion	/ /	:	/ /	:	/ /	:	1 1	:	
dates & times	/	<u> </u>	/ /	:	/_/_	<u>:</u>	/_/	:	

Go to the Discussions With Your Doctor page on ZYNLONTA.com for more information >









What is ZYNLONTA®?

ZYNLONTA® is a prescription medicine used to treat adults with certain types of large B-cell lymphoma that has come back (relapsed) or that did not respond to previous treatment (refractory), who have already received two or more treatments for their cancer.

It is not known if ZYNLONTA® is safe and effective in children.

The approval of ZYNLONTA® is based on a type of response rate. There is an ongoing study to confirm the clinical benefit of ZYNLONTA®.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of ZYNLONTA®?

ZYNLONTA® may cause serious side effects, including:

- Fluid retention. Your body may hold too much fluid during treatment with ZYNLONTA®. This can be serious. Tell your healthcare provider if you develop new or worsening swelling or puffiness, weight gain, chest pain, shortness of breath, or trouble breathing
- Low blood cell counts (platelets, red blood cells, and white blood cells). Low blood cell counts are common with ZYNLONTA® but can also be serious or severe. Your

healthcare provider will monitor your blood counts during treatment with ZYNLONTA®. Tell your healthcare provider right away if you get a fever of 100.4°F (38°C) or above, or any bruising or bleeding

- Infections. Serious infections, including infections that can cause death, have happened in people treated with ZYNLONTA®. Tell your healthcare provider right away if you have new or worsening signs or symptoms of infection
- Skin reactions. Serious skin reactions have happened in people treated with ZYNLONTA®. Tell your healthcare provider if you get new or worsening skin reactions, including sensitivity to sunlight, skin rash, peeling, redness or irritation. You may burn more easily or get severe sunburns

The most common side effects of ZYNLONTA® include:

- feeling tired or weak
- skin rash
- swelling
- nausea
- muscle or joint pain
- · increase in blood sugar (hyperglycemia)
- · changes in certain blood or laboratory tests







ZYNLONTA® may cause fertility problems in males which may affect your ability to father children. Talk to your healthcare provider if this is a concern for you. These are not all of the possible side effects of ZYNLONTA®.

What should I avoid while receiving ZYNLONTA®?

Avoid or limit your exposure to sunlight, including sunlight through glass, such as buildings or vehicle windows and artificial sunlight such as sunlamps or tanning beds. Exposure to sunlight during treatment with ZYNLONTA® can cause skin reaction or rash. Use sun protection measures such as sunscreen and wear loose-fitting clothes that cover your skin while out in sunlight.

Before you receive ZYNLONTA®, tell your healthcare provider about all of your medical conditions, including if you:

- have an active infection or have had one recently
- have liver problems
- are pregnant or plan to become pregnant. ZYNLONTA® can harm your unborn baby

Females who can become pregnant:

 your healthcare provider may do a pregnancy test before starting treatment with ZYNLONTA® you should use effective birth control (contraception)
during treatment with ZYNLONTA® and for 10 months
after the last dose of ZYNLONTA®. Talk to your healthcare
provider about effective birth control. Tell your healthcare
provider right away if you become pregnant or think that
you are pregnant during treatment with ZYNLONTA®

Males with female partners who can become pregnant:

- you should use effective birth control (contraception) during treatment with ZYNLONTA® and for 7 months after the last dose of ZYNLONTA®
- are breastfeeding or plan to breastfeed. It is not known if ZYNLONTA® passes into breast milk. Do not breastfeed during treatment with ZYNLONTA® and for 3 months after the last dose of ZYNLONTA®

Tell your healthcare provider about all the medicines that you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get new medicine.

You may report side effects to the FDA at **(800) FDA-1088** or www.fda.gov/medwatch. You may also report side effects to ADC Therapeutics at **1-855-690-0340**.







Resources and tips to help you and the person you care for navigate DLBCL

We are happy to provide resources to help you best manage your treatment goals. Find information about:



Help with your copay





Driving the conversations with your doctor



How ZYNLONTA $^{\!\scriptscriptstyle{(\! g)}}$ may help you

Your "here" matters

Maybe it's an event, place, or just being around someone special. Your "here" is where you feel most like yourself. **ZYNLONTA**® may help you get there.

Visit the resources page on ZYNLONTA.com to find out more >



Please see Important Safety Information throughout this brochure, and full **Prescribing Information** for ZYNLONTA®, including Patient Information, on **ZYNLONTA.com**.

